

Please write clearly in block capitals.

Centre number

Candidate number

Surname _____

Forename(s) _____

Candidate signature _____

A-level FURTHER MATHEMATICS

Paper 3 Statistics

Thursday 13 June 2019

Afternoon

Time allowed: 2 hours

Materials

- You must have the AQA formulae and statistical tables booklet for A-level Mathematics and A-level Further Mathematics.
- You should have a scientific calculator that meets the requirements of the specification. (You may use a graphical calculator.)
- You must ensure you have the other optional Question Paper/Answer Book for which you are entered (**either** Discrete **or** Mechanics). You will have 2 hours to complete **both** papers.

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer each question in the space provided for that question. If you require extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do **not** write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work you do not want to be marked.

| For Examiner's Use | |
|--------------------|------|
| Question | Mark |
| 1 | |
| 2 | |
| 3 | |
| 4 | |
| 5 | |
| 6 | |
| 7 | |
| 8 | |
| TOTAL | |

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 50.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.



Answer **all** questions in the spaces provided.

1 The discrete random variable X has $\text{Var}(X) = 5$

Find $\text{Var}(4X - 3)$

Circle your answer.

[1 mark]

17

20

77

80

2 Amy takes a sample of size 50 from a normal distribution with mean μ and variance 16

She conducts a hypothesis test with hypotheses:

$$H_0 : \mu = 52$$

$$H_1 : \mu > 52$$

She rejects the null hypothesis if her sample has a mean greater than 53

The actual population mean is 53.5

Find the probability that Amy makes a Type II error.

Circle your answer.

[1 mark]

0.4%

3.9%

18.9%

45.0%



3 Alan's journey time to work can be modelled by a normal distribution with standard deviation 6 minutes.

Alan measures the journey time to work for a random sample of 5 journeys. The mean of the 5 journey times is 36 minutes.

3 (a) Construct a 95% confidence interval for Alan's mean journey time to work, giving your values to one decimal place.

[2 marks]

3 (b) Alan claims that his mean journey time to work is 30 minutes.

State, with a reason, whether or not the confidence interval found in part (a) supports Alan's claim.

[1 mark]

3 (c) Suppose that the standard deviation is not known but a sample standard deviation is found from Alan's sample and calculated to be 6

Explain how the working in part (a) would change.

[1 mark]

Turn over ►



4 (b) A 6 metre clothes line is connected between the point P on one building and the point Q on a second building.

Roy is concerned the clothes line may break. He uses the random variable X to model the distance in metres from P where the clothes line breaks.

4 (b) (i) State a criticism of Roy's model.

[1 mark]

4 (b) (ii) On the axes below, sketch the probability density function for an alternative model for the clothes line.

[1 mark]



Turn over for the next question

Turn over ►



- 5 (c)** The insurance company finds that the maximum possible claim is £2000 and they decide to refine their probability density function.

Suggest how this could be done.

[2 marks]

Turn over for the next question

Turn over ►



6 (b) Rebecca claims that if the result of the test in part **(a)** is to reject the null hypothesis then it is easier to pass a driving test at centre *B* than centre *A*.

State, with a reason, whether or not you agree with Rebecca's claim.

[1 mark]

Turn over for the next question

Turn over ►



*Do not write
outside the
box*

Turn over for the next question

Turn over ►



8 The number of telephone calls received by an office can be modelled by a Poisson distribution with mean 3 calls per 10 minutes.

8 (a) Find the probability that:

8 (a) (i) the office receives exactly 2 calls in 10 minutes;

[1 mark]

8 (a) (ii) the office receives more than 30 calls in an hour.

[3 marks]

8 (b) The office manager splits an hour into 6 periods of 10 minutes and records the number of telephone calls received in each of the 10 minute periods.

Find the probability that the office receives exactly 2 calls in a 10 minute period exactly twice within an hour.

[3 marks]



8 (c) The office has just received a call.

8 (c) (i) Find the probability that the next call is received more than 10 minutes later.

[3 marks]

8 (c) (ii) Mahah arrives at the office 5 minutes after the last call was received.

State the probability that the next call received by the office is received more than 10 minutes later.

Explain your answer.

[2 marks]

END OF QUESTIONS



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



