

## Harmonic identities

### Starter

1. Find an approximation of  $\frac{1 - \cos 2\theta}{\theta \tan \theta}$  when  $\theta$  is small.

**Working:** 
$$\frac{1 - \cos 2\theta}{\theta \tan \theta} \approx \frac{1 - (1 - \frac{1}{2}(2\theta)^2)}{\theta \times \theta} = \frac{2\theta^2}{\theta^2} = 2$$

2. Find  $\lim_{\theta \rightarrow 0} \frac{\sin 3\theta + \tan 5\theta}{2 \sin \theta}$ .

**Working:** 
$$\lim_{\theta \rightarrow 0} \frac{\sin 3\theta + \tan 5\theta}{2 \sin \theta} = \frac{3\theta + 5\theta}{2\theta} = 4$$

3. Let  $3 \sin \theta + 4 \cos \theta \equiv R \sin(\theta + \alpha)$ , where  $R > 0$  and  $0^\circ < \alpha < 90^\circ$ .
- (a) By using the compound angle identity for  $\sin(A + B)$ , find values for  $R \sin \alpha$  and  $R \cos \alpha$ .
- (b) Using your answers to (a) find the values of  $R$  and  $\alpha$ .

**Working:** (a)  $3 \sin \theta + 4 \cos \theta \equiv R \sin(\theta + \alpha)$   
 $R \sin(\theta + \alpha) = R(\sin \theta \cos \alpha + \cos \theta \sin \alpha)$   
 $3 \sin \theta + 4 \cos \theta \equiv R \sin \theta \cos \alpha + R \cos \theta \sin \alpha$   
 Equating coefficients:

$$\begin{array}{ll} \sin \theta: & R \cos \alpha = 3 \\ \cos \theta: & R \sin \alpha = 4 \end{array}$$

(b) **Using the equations:**  $R \sin \alpha = 4$  and  $R \cos \alpha = 3$   
**Squaring and adding:**  $R^2 \sin^2 \alpha + R^2 \cos^2 \alpha = 4^2 + 3^2$   
 $R^2(\sin^2 \alpha + \cos^2 \alpha) = 25$

Since  $\sin^2 \alpha + \cos^2 \alpha = 1$ :  $R^2 = 25$   
 Since  $R > 0$ :  $R = 5$

**Divide the equations:**  $\frac{R \sin \alpha}{R \cos \alpha} = \frac{4}{3}$   
 $\tan \alpha = \frac{4}{3}$

$$\alpha = \tan^{-1} \frac{4}{3} \approx 53.1^\circ$$

The value of  $R$  and  $\alpha$  are 5 and  $53.1^\circ$  (3 s.f.) respectively.

4. Let  $12 \cos \theta + 5 \sin \theta \equiv R \cos(\theta - \alpha)$ , where  $R > 0$  and  $0^\circ < \alpha < 90^\circ$ . Using similar working to question 2, find the values of  $R$  and  $\alpha$ .

**Working:**  $12 \cos \theta + 5 \sin \theta \equiv R \cos(\theta - \alpha)$   
 $R \cos(\theta + \alpha) = R(\cos \theta \cos \alpha + \sin \theta \sin \alpha)$   
 $12 \cos \theta + 5 \sin \theta \equiv R \cos \theta \cos \alpha + R \sin \theta \sin \alpha$   
 Equating coefficients:

$$\cos \theta: \quad R \cos \alpha = 12$$

$$\sin \theta: \quad R \sin \alpha = 5$$

**Squaring and adding:**  $R^2 \cos^2 \alpha + R^2 \sin^2 \alpha = 12^2 + 5^2$

$$R^2(\cos^2 \alpha + \sin^2 \alpha) = 169$$

Since  $\sin^2 \alpha + \cos^2 \alpha = 1$ :  $R^2 = 169$

Since  $R > 0$ :  $R = 13$

**Divide the equations:**  $\frac{R \sin \alpha}{R \cos \alpha} = \frac{5}{12}$

$$\tan \alpha = \frac{5}{12}$$

$$\alpha = \tan^{-1} \frac{5}{12} \approx 22.6^\circ$$

The value of  $R$  and  $\alpha$  are 13 and  $22.6^\circ$  (3 s.f.) respectively.

- E.g. 1** (a) Given that  $a \sin \theta + b \cos \theta \equiv R \sin(\theta + \alpha)$ , where  $R > 0$  and  $0^\circ < \alpha < 90^\circ$ , write down expressions for  $R$  and  $\alpha$  in terms of  $a$  and  $b$ .  
 (b) Given that  $a \cos \theta + b \sin \theta \equiv R \cos(\theta - \alpha)$ , where  $R > 0$  and  $0^\circ < \alpha < 90^\circ$ , write down expressions for  $R$  and  $\alpha$  in terms of  $a$  and  $b$ .

**Working:** (a)  $R = \sqrt{a^2 + b^2}$  and  $\alpha = \tan^{-1} \frac{b}{a}$

(b)  $R = \sqrt{a^2 + b^2}$  and  $\alpha = \tan^{-1} \frac{b}{a}$

- E.g. 2** (a) Let  $a \sin \theta - b \cos \theta \equiv R \sin(\theta - \alpha)$ , where  $R > 0$  and  $0^\circ < \alpha < 90^\circ$ .  
Find expressions for  $R$  and  $\alpha$  in terms of  $a$  and  $b$ .
- (b) Hence write down expressions for  $R$  and  $\alpha$  in terms of  $a$  and  $b$  given that  
 $a \cos \theta - b \sin \theta \equiv R \cos(\theta + \alpha)$ , where  $R > 0$  and  $0^\circ < \alpha < 90^\circ$ .

**Working:** (a)  $a \sin \theta - b \cos \theta \equiv R \sin(\theta - \alpha)$   
 $R \sin(\theta - \alpha) = R(\sin \theta \cos \alpha - \cos \theta \sin \alpha)$   
 $a \sin \theta - b \cos \theta \equiv R \sin \theta \cos \alpha - R \cos \theta \sin \alpha$   
 Equating coefficients:  
 $\sin \theta: \quad R \cos \alpha = a$   
 $\cos \theta: \quad R \sin \alpha = b$   
**Squaring and adding:**  $R^2 \sin^2 \alpha + R^2 \cos^2 \alpha = a^2 + b^2$   
 Since  $\sin^2 \alpha + \cos^2 \alpha = 1$ :  $R^2 = a^2 + b^2$   
 Since  $R > 0$ :  $R = \sqrt{a^2 + b^2}$   
**Divide the equations:**  $\frac{R \sin \alpha}{R \cos \alpha} = \frac{b}{a}$   
 $\tan \alpha = \frac{b}{a}$   
 $\alpha = \tan^{-1} \frac{b}{a}$

(b)  $R = \sqrt{a^2 + b^2}$  and  $\alpha = \tan^{-1} \frac{b}{a}$

- E.g. 3** (a) Write  $2 \sin \theta - 9 \cos \theta$  in the form  $R \sin(\theta - \alpha)$ , where  $R > 0$  and  $0^\circ < \alpha < 90^\circ$ .
- (b) Write  $7 \cos \theta - 3 \sin \theta$  in the form  $R \cos(\theta + \alpha)$ , where  $R > 0$  and  $0^\circ < \alpha < 90^\circ$ .

**Working:** (a)  $R = \sqrt{2^2 + 9^2} = \sqrt{85}$  and  $\alpha = \tan^{-1} \left| \frac{9}{2} \right| = 77.5^\circ$   
 $2 \sin \theta - 9 \cos \theta \equiv \sqrt{85} \sin(\theta - 77.5^\circ)$

(b)  $R = \sqrt{7^2 + 3^2} = \sqrt{58}$  and  $\alpha = \tan^{-1} \left| \frac{3}{7} \right| = 23.2^\circ$   
 $7 \cos \theta - 3 \sin \theta \equiv \sqrt{58} \cos(\theta + 23.2^\circ)$

- E.g. 4** (a) Solve the equation  $\sin \theta + \sqrt{3} \cos \theta = 1$  for  $0^\circ \leq \theta \leq 360^\circ$ .  
 (b) Solve the equation  $4 \cos \theta + 9 \sin \theta = 5$  for  $0^\circ \leq \theta \leq 2\pi$ .

**Working:**

(a) Let  $\sin \theta + \sqrt{3} \cos \theta \equiv R \sin(\theta + \alpha)$   
 $R = \sqrt{1^2 + 3} = 2$  and  $\alpha = \tan^{-1} \left| \frac{\sqrt{3}}{1} \right| = 60^\circ$   
 $\sin \theta + \sqrt{3} \cos \theta \equiv 2 \sin(\theta + 60^\circ)$   
 $\sin \theta + \sqrt{3} \cos \theta = 1 \Rightarrow 2 \sin(\theta + 60^\circ) = 1$   
 $\sin(\theta + 60^\circ) = \frac{1}{2}$   
 Since  $\sin^{-1} \frac{1}{2} = 30^\circ$  and  $\frac{1}{2}$  is positive:  
 $\theta + 60^\circ = 30^\circ$  (A quadrant)  $\Rightarrow \theta = -30^\circ$   
 or  $\theta + 60^\circ = 180^\circ - 30^\circ$  (S quadrant)  $\Rightarrow \theta = 90^\circ$   
 Since  $-30^\circ$  is not  $0^\circ \leq \theta \leq 360^\circ$ , add  $360^\circ$ .  
 So  $\theta = 90^\circ$  and  $\theta = 330^\circ$

(b) Let  $4 \cos \theta + 9 \sin \theta \equiv R \cos(\theta - \alpha)$   
 $R = \sqrt{4^2 + 9^2} = \sqrt{97}$  and  $\alpha = \tan^{-1} \left| \frac{9}{4} \right| = 1.1526^c$   
 $4 \cos \theta + 9 \sin \theta \equiv \sqrt{97} \cos(\theta - 1.1526^c)$   
 $4 \cos \theta + 9 \sin \theta = 5 \Rightarrow \sqrt{97} \cos(\theta - 1.1526^c) = 5$   
 $\cos(\theta - 1.1526^c) = \frac{5}{\sqrt{97}}$   
 Since  $\cos^{-1} \frac{5}{\sqrt{97}} = 1.0383^c$  and  $\frac{5}{\sqrt{97}}$  is positive:  
 $\theta - 1.1526^c = 1.0383^c$  (A quadrant)  $\Rightarrow \theta = 2.19^c$   
 or  $\theta - 1.1526^c = 2\pi - 1.0383^c$  (C quadrant)  $\Rightarrow \theta = 6.40^c$   
 Since  $6.40^c$  is not  $0^\circ \leq \theta \leq 2\pi$ , add  $2\pi$ .  
 So  $\theta = 0.114^c$  and  $\theta = 2.19^c$ .

- E.g. 5** Find the range of values of the constant  $k$  for which the equation  $\sin \theta + \sqrt{2} \cos \theta = k$  has real solutions for  $\theta$ .

**Working:** Let  $\sin \theta + \sqrt{2} \cos \theta \equiv R \sin(\theta - \alpha)$  then  $R = \sqrt{1^2 + 2} = \sqrt{3}$ .  
 $\sin \theta + \sqrt{2} \cos \theta = k \Rightarrow \sqrt{3} \sin(\theta + \alpha) = k$   
 $\sin(\theta + \alpha) = \frac{k}{\sqrt{3}}$   
 Since  $-1 \leq \sin x \leq 1$ :  $-1 \leq \frac{k}{\sqrt{3}} \leq 1$   
 Multiply by  $\sqrt{3}$ :  $-\sqrt{3} \leq k \leq \sqrt{3}$

**E.g. 6** State the transformations that take the curve  $y = \cos x$  to the curve  $y = 8 \cos \theta - 3 \sin \theta$ .

**Working:** Let  $8 \cos \theta - 3 \sin \theta \equiv R \cos(\theta + \alpha)$

$$R = \sqrt{8^2 + 3^2} = \sqrt{73} \quad \text{and} \quad \alpha = \tan^{-1} \left| \frac{3}{8} \right| = 20.6^\circ$$

$$y = 8 \cos \theta - 3 \sin \theta \quad \Rightarrow \quad y = \sqrt{73} \cos(\theta + 20.6^\circ)$$

The transformations are:

a stretch, factor  $\sqrt{73}$ , parallel to the  $y$ -axis.

a translation parallel to the  $x$ -axis,  $20.6^\circ$  (or  $0.359^c$ ) units to the left

**E.g. 7** (a) Express  $f(\theta) = 7 \sin \theta - 24 \cos \theta$  in the form  $R \sin(\theta - \alpha)$ , where  $R > 0$  and  $0^\circ < \alpha < 90^\circ$ .

(b) Hence find the greatest value of  $\frac{9}{f(\theta) + 40}$ . State the value of  $\theta$  for which this occurs.

**Working:** (a)  $R = \sqrt{7^2 + 24^2} = 25$  and  $\alpha = \tan^{-1} \left| \frac{24}{7} \right| = 73.7^\circ$   
 $f(\theta) = 25 \sin(\theta - 73.7^\circ)$

$$(b) \quad \frac{9}{f(\theta) + 40} = \frac{9}{25 \sin(\theta - 73.7^\circ) + 40}$$

The greatest value is when the denominator is the least value.

$$\text{Since } -1 \leq \sin x \leq 1: \quad -25 \leq 25 \sin(\theta - 73.7^\circ) \leq 25$$

$$\text{Adding 40:} \quad 15 \leq 25 \sin(\theta - 73.7^\circ) + 40 \leq 65.$$

The greatest value is when the denominator is the least.

$$\text{The greatest value of } \frac{9}{f(\theta) + 40} \text{ is } \frac{9}{-25 + 40} = \frac{9}{15} = \frac{3}{5}.$$

$$\text{This value occurs when } \sin(\theta - 73.7^\circ) = -1$$

$$\theta - 73.7^\circ = 270^\circ$$

$$\theta = 344^\circ \text{ (3 s.f.)}$$

**Video:** [Harmonic identities](#)

[Harmonic identities EQ](#)

[Solutions to Starter and E.g.s](#)

### Exercise

p177 8C Qu 1-8, (9-10 red)