

Modelling Projectile Motion

Starter

1. **(Review of AS material)** A particle is projected vertically upward from ground level with a speed of 24.5 m/s. Find:
- the greatest height reached
 - the time that elapses before the particle returns to the ground.

Working:

(a) At the greatest height $v = 0$
 So $u = 24.5, v = 0, a = -9.8, s = ?$
 No $t \Rightarrow v^2 = u^2 + 2as: 0 = 24.5^2 + 2 \times -9.8 \times s$
 $s = 30.6$ so the greatest height is 30.6 m

(b) When the particle returns to ground $s = 0$
 So $u = 24.5, a = -9.8, s = 0, t = ?$
 No $v \Rightarrow s = ut + \frac{1}{2}at^2: 0 = 24.5t + \frac{1}{2} \times (-9.8) \times t^2$
 $t(24.5 - 4.9t) = 0 \Rightarrow t = 0$ or $t = \frac{24.5}{4.9} = 5$
 \therefore the time to return to ground is 5 seconds

2. **(Review of AS material)** A particle starts from the origin and moves under acceleration $\mathbf{a} = (2\mathbf{i} - 7\mathbf{j})$ m/s². Given that the initial velocity is $\mathbf{u} = (-\mathbf{i} + 6\mathbf{j})$. Find:
- the position and velocity of the particle after 4 s and
 - the time when the particle is at the point $(\lambda\mathbf{i} - 12\mathbf{j})$, where λ is a value to be found.

Working:

(a) Position: $\mathbf{s} = \mathbf{u}t + \frac{1}{2}\mathbf{a}t^2$
 $\mathbf{s} = (-\mathbf{i} + 6\mathbf{j}) \times 4 + 0.5 \times (2\mathbf{i} - 7\mathbf{j}) \times 4^2$
 $\mathbf{s} = (12\mathbf{i} - 32\mathbf{j})$ m

Velocity: $\mathbf{v} = \mathbf{u} + \mathbf{a}t$
 $\mathbf{v} = (-\mathbf{i} + 6\mathbf{j}) + (2\mathbf{i} - 7\mathbf{j}) \times 4$
 $\mathbf{v} = (7\mathbf{i} - 22\mathbf{j})$ m/s

(b) $s = ut + \frac{1}{2}at^2$
 $\mathbf{j}: -12 = 6t + 0.5 \times (-7) \times t^2$
 $t = 2.90$ or $t = -1.18$
 $\mathbf{i}: \text{Since } t > 0, \text{ when } t = 2.90,$
 $\lambda = -1 \times 2.90 + 0.5 \times 2 \times 2.90^2 = 5.51$

E.g. 1 Find the horizontal and vertical components of the velocity when a particle is moving with speed 8 m/s at an angle of 35° above the horizontal.

Working: Horizontal = $8 \cos 35 = 6.55$ m/s; *turn through 35° so cos*
 Vertical = $8 \sin 35 = 4.59$ m/s *not turn through 35° so sin*

E.g. 2 A ball is thrown with velocity v , with horizontal component 6 m/s and vertical component 8 m/s. Find the speed and direction of projection of the ball.

Working: Speed = $\sqrt{6^2 + 8^2} = 10$ m/s;
Direction = $\tan^{-1} \frac{8}{6} = 53.1^\circ$ above the horizontal

E.g. 3 A particle is projected with initial speed 18 m/s at an angle of 40° above the horizontal. Find the distance from its point of projection 1.8 s after it is projected.

Working: *The aim is to find the distance travelled horizontally and vertically and then use Pythagoras.*

$$u_x = 18 \cos 40, a_x = 0, t = 1.8, s_x = ? \quad (\text{horizontally})$$

$$\text{No } v_x \Rightarrow s = ut + \frac{1}{2}at^2$$
$$s_x = 18 \times 1.8 \times \cos 40 + 0 = 24.8\dots$$

$$u_y = 18 \sin 40, a_y = -9.8, t = 1.8, s_y = ? \quad (\text{vertically})$$

$$\text{No } v_y \Rightarrow s = ut + \frac{1}{2}at^2$$
$$s_y = 18 \times 1.8 \times \sin 40 - 0.5 \times 9.8 \times 1.8^2 = 4.95\dots$$

$$\text{Distance} = \sqrt{24.8^2 + 4.95^2} = 25.3 \text{ m (3 s.f.)}$$

E.g. 4 A particle is projected from the point $5\mathbf{j}$. The particle's initial velocity is $(7\mathbf{i} + 3\mathbf{j})$ m/s and it moves freely under gravity.

- (a) Find its position vector 0.8 s after it is projected.
(b) Find the velocity at this time.

Working: (a) $\mathbf{u} = 7\mathbf{i} + 3\mathbf{j}, t = 0.8, \mathbf{a} = -9.8\mathbf{j}, \mathbf{s} = ?$

$$\text{No } \mathbf{v} \Rightarrow \mathbf{s} = \mathbf{u}t + \frac{1}{2}\mathbf{a}t^2$$

The starting position is $5\mathbf{j}$ so:

$$\mathbf{s} = 5\mathbf{j} + (7\mathbf{i} + 3\mathbf{j}) \times 0.8 + \frac{1}{2} \times -9.8\mathbf{j} \times 0.8^2$$
$$= (5.6\mathbf{i} - 4.264\mathbf{j})$$

The position vector after 0.8 s is $(5.6\mathbf{i} + 4.264\mathbf{j})\text{m}$

(b) $\mathbf{u} = 7\mathbf{i} + 3\mathbf{j}, t = 0.8, \mathbf{a} = -9.8\mathbf{j}, \mathbf{v} = ?$

$$\text{No } \mathbf{s} \Rightarrow \mathbf{v} = \mathbf{u} + \mathbf{a}t$$

$$\mathbf{v} = (7\mathbf{i} + 3\mathbf{j}) + (0\mathbf{j} - 9.8\mathbf{j}) \times 0.8 = (7\mathbf{i} - 4.84\mathbf{j}) \text{ m/s}$$

[Video: Projectiles](#)

[Video: Projectiles \(upward angle/above ground\)](#)

[Video: Projectiles \(downward angle/above ground\)](#)

[Video: Projected horizontally](#)

[Solutions to Starter and E.g.s](#)

Exercise

p455 20A Qu 1-4

Answers to additional questions

1. Horizontal = $6.11 \cos \alpha$ m/s; vertical = $6.11 \sin \alpha$ m/s
2. (a) $v_x = 10, v_y = -4.9$ so speed = 11.1 m/s (3 s.f.)
and direction = 26.1° (3 s.f.) below horizontal
(b) $s_y = -2, t = 0.6388\dots$ so $s_x = 6.39$
3. (a) $(12\mathbf{i} - 3.6\mathbf{j})$ m/s
(b) $12\mathbf{i}$ m/s
(c) $(12\mathbf{i} - 16\mathbf{j})$ m/s
4. (a) 10.1 m
(b) 22.1 m/s
(c) 39.6° below the horizontal
5. (a) 0.719 s
(b) 7.51 m
6. (a) $a = 40, b = 16.5$
(b) $\mathbf{v} = (40\mathbf{i} - 32.5\mathbf{j})$ m/s