


Non-uniform rods, tilting and moment of angled force

Starter

1. (Review of last lesson)

A uniform rod, AB , of length 5 m and mass M kg, rests horizontally in equilibrium on supports at C and D , as shown. If the magnitude of the normal reaction at C is 49 N, find:



- (a) the magnitude of the normal reaction at D ,
- (b) the mass, M , of the rod.

Working: (a) An unknown force acts through the centre of the rod so take moments about the centre.

$$\curvearrowleft \text{ about centre: } R_D \times 0.5 = 49 \times 1$$

$$R_D = 98 \text{ N}$$

The normal reaction at D is 98 N

(b) $R(\uparrow)$: $R_D + 49 = Mg$

$$98 + 49 = Mg$$


$$M = 15 \text{ kg}$$

The mass, M , of the rod is 15 kg

E.g. 1 A non-uniform beam, AB , of mass 3 kg and length 1.6 m rests horizontally in equilibrium on vertical supports at A and B . The normal reaction at A is 12 N. Find

- (a) the magnitude of the normal reaction at B
- (b) the distance of the centre of mass of the beam from A .

Working: (a) $R(\uparrow)$: $R_B + 12 = 3g$
 $R_B = 3g - 12$
 $R_B = 17.4 \text{ N}$



The magnitude of the normal reaction at B is 17.4 N.

(b) In case our answer to R_B is wrong, take moments about B .

$$\curvearrowleft \text{ about B: } 3g \times (1.6 - d) = 12 \times 1.6$$

$$4.8g - 3dg = 19.2$$


$$4.8g - 19.2 = 3dg$$

$$d = \frac{4.8g - 19.2}{3g}$$

$$d = \frac{232}{245} = 0.947 \text{ m (3 s.f.)}$$


The distance of the centre of mass of the beam from A is 0.947 m

- E.g. 2** A non-uniform rod of mass 1.5 kg rests horizontally in equilibrium on supports at A and B , as shown. When a downwards force of magnitude F N is applied to the rod at C , the rod remains horizontal and in equilibrium, but is on the point of tilting about B , where the normal reaction has magnitude 44 N. Given that the distance between B and the centre of mass is 2.5 m, find:
- the value of F ,
 - the distance between the points B and C .



- Working:**
- The rod is on the point of tilting about B so the normal reaction at A is zero i.e. $R_A = 0$
 $R(\uparrow): F + 1.5g = 44$
 $F = 44 - 1.5g$
 $F = 29.3$ N
 - Let d be the distance between B and C
 \curvearrowright about $B: 29.3 \times d = 1.5g \times 2.5$
 $d = \frac{735}{586} \approx 1.25$
 The distance between the points B and C is 1.25 m (3 s.f.)

- E.g. 3** A light rod, AB , has length 4 m. A force of magnitude 5 N is applied to A at an angle of 40° above the rod, as shown. A second force of magnitude 3 N is applied vertically upwards at the rod's midpoint. Find the sum of the moments of the forces acting on the rod about B .



- Working:** Resolve the 5 N force perpendicular to the rod: $5 \sin 40$
 \curvearrowright about $B: 5 \sin 40 \times 4 + 3 \times 2 \approx 18.9$
 The sum of the moments about B is 18.9 Nm (3 s.f.) clockwise.

Video: [Tilting](#)
 Video: [Moment of non-perpendicular force](#)

[Tilting EQ](#)
[Moment of non-perpendicular force EQ](#)

[Solutions to Starter and E.g.s](#)

Exercise
 p517 22C Qu 1-7