

## Radians

1. How many degrees are equivalent to one radian?

**Hint:** Arc length,  $s = \frac{\theta}{360^\circ} \times 2\pi r$

**Working:** From GCSE, arc length:  $s = \frac{\theta}{360^\circ} \times 2\pi r$   
 Since the angle is 1 radian:  $s = r$  (i.e. arc length = radius)  
 $r = \frac{\theta}{360^\circ} \times 2\pi r$   
 Cancel  $r$  each side:  $1 = \frac{2\pi\theta}{360^\circ}$   
 Rearranging gives:  $\theta = \frac{360^\circ}{2\pi} = \frac{180^\circ}{\pi} \approx 57.3^\circ$

1 radian  $\approx 57.3^\circ$

**E.g. 1** Convert  $45^\circ$  to radians.

**Working:**  $45^\circ \equiv 45 \times \frac{\pi}{180} = \frac{\pi}{4}$

Notice that since we have a fraction, we leave the angle in terms of  $\pi$ .

**E.g. 2** Convert the following to degrees: (a)  $\frac{2\pi}{3}$  (b)  $1.85^c$

**Working:** (a)  $\frac{2\pi}{3} \equiv \frac{2\pi}{3} \times \frac{180}{\pi} = 120^\circ$

(b)  $1.85^c \equiv 1.85^c \times \frac{180}{\pi} \approx 106^\circ$

### Common angles

**E.g. 3** Copy and complete this table.

<b>Degree</b>	$0^\circ$	$30^\circ$	$45^\circ$		$90^\circ$	$120^\circ$		$150^\circ$	$180^\circ$		$360^\circ$
<b>Radians</b>	$0^c$		$\frac{\pi}{4}$	$\frac{\pi}{3}$		$\frac{2\pi}{3}$	$\frac{3\pi}{4}$		$\pi$	$\frac{3\pi}{2}$	$2\pi$

**Working:**

<b>Degree</b>	$0^\circ$	$30^\circ$	$45^\circ$	$60^\circ$	$90^\circ$	$120^\circ$	$135^\circ$	$150^\circ$	$180^\circ$	$270^\circ$	$360^\circ$
<b>Radians</b>	$0^c$	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	$\pi$	$\frac{3\pi}{2}$	$2\pi$

**E.g. 4** Given that  $\sin \frac{\pi}{6} = \frac{1}{2}$  state the values of the following without using a calculator:

- (a)  $\sin \frac{13\pi}{6}$  (b)  $\sin \left( -\frac{\pi}{6} \right)$   
(c)  $\sin \frac{37\pi}{6}$  (d)  $\sin \frac{11\pi}{6}$

**Working:**

(a) Periodic:  $\sin \frac{13\pi}{6} = \sin \left( \frac{\pi}{6} + 2\pi \right) = \sin \frac{\pi}{6} = \frac{1}{2}$

(b) Symmetry:  $\sin \left( -\frac{\pi}{6} \right) = -\sin \left( \frac{\pi}{6} \right) = -\frac{1}{2}$

(c) Periodic:  $\sin \frac{37\pi}{6} = \sin \left( \frac{\pi}{6} + 6\pi \right) = \sin \frac{\pi}{6} = \frac{1}{2}$

(d) Combining:  $\sin \frac{11\pi}{6} = \sin \left( 2\pi - \frac{\pi}{6} \right) = -\sin \frac{\pi}{6} = -\frac{1}{2}$

**E.g. 5** Given that  $\cos \frac{\pi}{6} = \frac{\sqrt{3}}{2}$  state the values of the following without using a calculator:

- (a)  $\cos \frac{25\pi}{6}$  (b)  $\cos \left( -\frac{\pi}{6} \right)$   
(c)  $\cos \left( -\frac{11\pi}{6} \right)$  (d)  $\cos \frac{7\pi}{6}$

**Working:**

(a)  $\cos \frac{25\pi}{6} = \cos \left( \frac{25\pi}{6} - 4\pi \right) = \cos \frac{\pi}{6} = \frac{\sqrt{3}}{2}$

(b)  $\cos \left( -\frac{\pi}{6} \right) = \cos \frac{\pi}{6} = \frac{\sqrt{3}}{2}$

(c)  $\cos \left( -\frac{11\pi}{6} \right) = \cos \left( -\frac{11\pi}{6} + 2\pi \right) = \cos \frac{\pi}{6} = \frac{\sqrt{3}}{2}$

(d)  $\cos \frac{7\pi}{6} = \cos \left( \frac{7\pi}{6} - 2\pi \right) = \cos \left( -\frac{5\pi}{6} \right)$   
 $= -\cos \left( \pi - \frac{5\pi}{6} \right) = -\cos \frac{\pi}{6} = -\frac{\sqrt{3}}{2}$

**Video: Radians**

**Solutions to Starter and E.g.s**

**Exercise**

p133 7A Qu 2i, 3i, 4i, 6-12