

Please write clearly, in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

Forename(s)

Candidate signature

AS FURTHER MATHEMATICS

Paper 2 – Mechanics

Exam Date

Morning

Time allowed: 1 hour 30 minutes

Materials

For this paper you must have:

- You must ensure you have the other optional question paper/answer booklet for which you are entered (**either** Discrete **or** Statistics). You will have 1 hour 30 minutes to complete both papers.
- The AQA booklet of formulae and statistical tables.
- You may use a graphics calculator.

Instructions

- Use black ink or black ball-point pen. Pencil should be used for drawing.
- Answer **all** questions.
- You must answer each question in the space provided for that question. If you require extra space, use an AQA supplementary answer book; do **not** use the space provided for a different question.
- Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 40.

Advice

Unless stated otherwise, you may quote formulae, without proof, from the booklet.
You do not necessarily need to use all the space provided.

Answer **all** questions in the spaces provided.

1 A child, of mass 40 kg, moves at constant speed of 5 m s^{-1} on a fairground ride.

The path of the child is a circle of radius 4 metres.

Find the magnitude of the resultant force acting on the child.

Circle your answer.

[1 mark]


6.3 N

50 N

130 N

250 N

- 2 The graph shows how a force, F , varies with time during a period of 0.8 seconds.



Find the magnitude of the impulse of F during the 0.8 seconds.

Circle your answer.

[1 mark]

1.0 Ns

1.6 Ns

2.2 Ns

3.2 Ns

Turn over for the next question

3 A tank full of liquid has a hole made in its base.

Two students, Sarah and David, propose two different models for the speed, v , at which liquid exits the tank.

David thinks that v will depend on the height of the liquid in the tank, h , the acceleration due to gravity, g , and the density of the liquid, ρ , such that $v \propto g^a h^b \rho^c$ where a , b and c are constants.

Sarah thinks that v will not depend on the density of the liquid and suggests the model $v \propto g^a h^b$

3 (a) By considering dimensions, explain which student's model should be rejected.

[2 marks]

3 (b) Find the values of the constants in order for the model that you did **not** reject in part (a) to be dimensionally consistent.

[2 marks]

5 (b) Find the speed of C immediately after the collision.

[2 marks]

5 (c) In fact the horizontal surface on which the discs are sliding is not smooth.
Explain how the introduction of friction will affect your answer to part **(b)**.

[2 marks]

Turn over for the next question

Turn over ▶

- 7 A disc, of mass 0.15 kg, slides across a smooth horizontal table and collides with a vertical wall which is perpendicular to the path of the disc.

The disc is in contact with the wall for 0.02 seconds and then rebounds.

A possible model for the force, F newtons, exerted on the disc by the wall, whilst in contact, is given by

$$F = kt^2(t-b)^2 \quad \text{for } 0 \leq t \leq 0.020$$

where k and b are constants.

The force is initially zero and becomes zero again as the disc loses contact with the wall.

- 7 (a) State the value of b .

[1 mark]

- 7 (b) Find the magnitude of the impulse on the disc, giving your answer in terms of k .

[3 marks]

-
- 7 (c)** The disc is travelling at 4 m s^{-1} when it hits the wall.
The disc rebounds with a speed of 2 m s^{-1}
Find k .

[3 marks]

Turn over for the next question

8 (b) Describe one way in which the model you have used could be refined.

[1 mark]

END OF QUESTIONS