

Collisions and the principle of conservation of momentum

Starter

1. **(Review of last lesson)** A particle of mass 2 kg is moving in a straight line, with a speed of 5 m/s. A force of 11 N acts on the particle for 6 seconds, in the direction of motion. Find
- the magnitude of the change in momentum of the particle
 - the speed of the particle at the end of this time.

Working: (a) Change in momentum = $Ft = 11 \times 6 = 66$ Ns

$$(b) \quad \text{Increase in speed } Ft = mv - mu: \quad \begin{aligned} 66 &= 2v - 2 \times 5 \\ 2v &= 76 \\ v &= 38 \end{aligned}$$

The speed of the particle at the end of this time is 38 m/s.

2. **(Review of last lesson)** A body of mass 5 kg is moving with speed 7 m/s when a force is applied to it for 8 seconds. Its speed then is again 7 m/s but in the opposite direction. Find the magnitude of the force that has caused this change.

Working: $Ft = mu - mv: \quad \begin{aligned} F \times 8 &= 5 \times 7 - 5 \times (-7) \\ F &= \frac{35 - -35}{8} \\ F &= 8.75 \end{aligned}$

The magnitude of the force is 8.75 N.


3. An object A of mass m_A collides with object B which has mass m_B . Prior to the collision they had velocities u_A and u_B respectively and after the collision their velocities are v_A and v_B . Find the impulse of both objects.

Working: Impulse of object A = change in momentum = $m_A v_A - m_A u_A$.
Impulse of object B = change in momentum = $m_B v_B - m_B u_B$.

- E.g. 1** A particle, A , of mass 3 kg, travelling at 5 m/s collides head-on with a particle B with mass 2 kg and travelling at 4 m/s. If, after impact, B moves in the opposite direction at 2 m/s, find the velocity of A .

N.B. A "head-on" collision means the objects are moving towards each other.

Working:



$$\text{CoM:} \quad 3 \times 5 - 2 \times 4 = 3v_A + 2 \times 2$$

Note the negative sign since B is initially travelling in the opposite direction.

$$7 = 3v_A + 4$$


$$v_A = 1$$

Since $v_A > 0$ it is travelling in the defined positive direction.

The velocity of A is 1 m/s in its original direction of motion.

E.g. 2 A bullet of mass 0.1 kg is fired horizontally, at 80 m/s, into a stationary block of wood that is free to move on a smooth horizontal plane. The wooden block, with the bullet embedded in it, moves off with speed 5 m/s. Find the mass of the block.

Working:




$$\begin{aligned} \text{CoM:} \quad 0.1 \times 80 &= (m + 0.1) \times 5 \\ 8 &= 5m + 0.5 \\ m &= 1.5 \end{aligned}$$

The mass of the block is 1.5 kg

E.g. 3 A particle, *A*, of mass 5 kg travelling with speed 6 m/s, collides directly with a stationary particle *B* of mass 10 kg. If *A* is brought to rest by the impact find the speed with which *B* begins to move.

Working:



$$\begin{aligned} \text{CoM:} \quad 5 \times 6 &= 10 \times v_B \\ v_B &= 3 \end{aligned}$$


B begins to move with speed 3 m/s

E.g. 4 A three-tonne articulated lorry is moving along a track at 8 m/s towards a five-tonne bin lorry travelling at 5 m/s on the same road. If the lorries become coupled at impact find the velocity at which they continue to move if they were travelling

- (a) in the same direction
 (b) in the opposite direction.

Make sure you give the direction of motion.


Working: (a)



$$\begin{aligned} \text{CoM:} \quad 3000 \times 8 + 5000 \times 5 &= (3000 + 5000) \times v \\ 49000 &= 8000v \\ v &= 6.125 \end{aligned}$$

The velocity at which they continue to move is 6.75 m/s in the same direction as the original direction of the trucks.

(b)



$$\begin{aligned} \text{CoM:} \quad 3000 \times 8 - 5000 \times 5 &= (3000 + 5000) \times v \\ -1000 &= 8000v \\ v &= -0.125 \end{aligned}$$

The velocity at which they continue to move is 0.125 m/s in the same direction as the original direction of the bin lorry.

Video: [Conservation of linear momentum](#)

Video: [Colliding and separating](#)

[Solutions to Starter and E.g.s](#)

Exercise

p60 3B Qu 1-10