

Vector Geometry

Starter

1. (Review of last lesson)

The position vectors of 3 vertices of a parallelogram are $\begin{pmatrix} 4 \\ 2 \end{pmatrix}$, $\begin{pmatrix} 3 \\ 5 \end{pmatrix}$ and $\begin{pmatrix} 8 \\ 6 \end{pmatrix}$. Find the two possible position vectors of the 4th vertex in the first quadrant.

Working: Draw a quick diagram and label the points to help.


$$\vec{BC} = \begin{pmatrix} 8 \\ 6 \end{pmatrix} - \begin{pmatrix} 3 \\ 5 \end{pmatrix} = \begin{pmatrix} 5 \\ 1 \end{pmatrix}$$

$$\text{So 4th vertex} = \begin{pmatrix} 4 \\ 2 \end{pmatrix} + \begin{pmatrix} 5 \\ 1 \end{pmatrix} = \begin{pmatrix} 9 \\ 3 \end{pmatrix}$$

$$\vec{AC} = \begin{pmatrix} 8 \\ 6 \end{pmatrix} - \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$\text{So 4th vertex} = \begin{pmatrix} 3 \\ 5 \end{pmatrix} + \begin{pmatrix} 4 \\ 4 \end{pmatrix} = \begin{pmatrix} 7 \\ 9 \end{pmatrix}$$

$$\text{4th vertex is } \begin{pmatrix} 7 \\ 9 \end{pmatrix} \text{ or } \begin{pmatrix} 9 \\ 3 \end{pmatrix}$$



E.g. 1 A , B and C are the points $(2, 5)$, $(4, 9)$ and $(-3, -5)$.

- (a) Find the vectors \vec{AB} and \vec{BC} .
 (b) Show that all three points are collinear.

Working: (a) $\vec{AB} = \begin{pmatrix} 4 \\ 9 \end{pmatrix} - \begin{pmatrix} 2 \\ 5 \end{pmatrix} = \begin{pmatrix} 2 \\ 4 \end{pmatrix}$

$$\vec{BC} = \begin{pmatrix} -3 \\ -5 \end{pmatrix} - \begin{pmatrix} 4 \\ 9 \end{pmatrix} = \begin{pmatrix} -7 \\ -14 \end{pmatrix}$$

(b) $\vec{BC} = -\frac{7}{2}\vec{AB}$ therefore the vectors are multiples of one another.

This means that \vec{AB} and \vec{BC} are parallel.

Since both vectors pass through B , the points are collinear.

E.g. 2 M is the midpoint of the line PQ , where P has position vector $-3\mathbf{i} + \mathbf{j}$ and M has position vector $2\mathbf{i} - 5\mathbf{j}$. What is the position vector of Q ?

Hint: if you are not sure what to do, draw a diagram.

Working: Since $\vec{PM} = \vec{MQ}$, we can find the position vector of Q by adding \vec{MQ} to the position vector of M

$$\vec{PM} = \begin{pmatrix} 2 \\ -5 \end{pmatrix} - \begin{pmatrix} -3 \\ 1 \end{pmatrix} = \begin{pmatrix} 5 \\ -6 \end{pmatrix} \equiv \vec{MQ}$$

$$\text{Position vector of } Q = \begin{pmatrix} 2 \\ -5 \end{pmatrix} + \begin{pmatrix} 5 \\ -6 \end{pmatrix} = \begin{pmatrix} 7 \\ -11 \end{pmatrix} \quad \text{i.e. } 7\mathbf{i} - 11\mathbf{j}$$

E.g. 3 Let $\vec{OA} = \mathbf{a}$ and let T be on OA such that T divides OA in the ratio 3 : 1.

(a) Is T closer to O or closer to A?

(b) Express \vec{OT} in terms of \mathbf{a} .


(c) Express \vec{AT} in terms of \mathbf{a} .

Hint: Draw a diagram.

Working: (a) Closer to A

(b) $\vec{OT} = \frac{3}{4}\mathbf{a}$

(c) $\vec{AT} = -\vec{TA} = -\frac{1}{4}\vec{OA} = -\frac{1}{4}\mathbf{a}$




E.g. 4 $\vec{AB} = \mathbf{p}$ and $\vec{AC} = \mathbf{q}$. The point X lies on BC and divides it in the ratio 2 : 5. Find \vec{AX} in terms of \mathbf{p} and \mathbf{q} .

Hint: Draw a diagram.

Working: $\vec{AX} = \vec{AB} + \vec{BX} = \vec{AB} + \frac{2}{7}\vec{BC}$

Since $\vec{BC} = \mathbf{q} - \mathbf{p}$

$\vec{AX} = \mathbf{p} + \frac{2}{7}(\mathbf{q} - \mathbf{p}) = \frac{5}{7}\mathbf{p} + \frac{2}{7}\mathbf{q}$



E.g. 5 Four points have coordinates A(2, -1), B(k, k + 1), C(2k - 3, 2k + 2) and D(k - 1, k).

(a) Show that ABCD is a parallelogram for all values of k.


(b) Find the value of k for which ABCD is a rhombus.

Working: (a) For a parallelogram we need to show $\vec{AB} = \vec{DC}$.

$\vec{AB} = \begin{pmatrix} k \\ k+1 \end{pmatrix} - \begin{pmatrix} 2 \\ -1 \end{pmatrix} = \begin{pmatrix} k-2 \\ k+2 \end{pmatrix}$

$\vec{DC} = \begin{pmatrix} 2k-3 \\ 2k+2 \end{pmatrix} - \begin{pmatrix} k-1 \\ k \end{pmatrix} = \begin{pmatrix} k-2 \\ k+2 \end{pmatrix}$

Since both \vec{AB} and \vec{DC} are equal to $\begin{pmatrix} k-2 \\ k+2 \end{pmatrix}$, ABCD is a parallelogram.



(b) For ABCD to be a rhombus $|\vec{AB}| = |\vec{AD}|$

$\vec{AD} = \begin{pmatrix} k-1 \\ k \end{pmatrix} - \begin{pmatrix} 2 \\ -1 \end{pmatrix} = \begin{pmatrix} k-3 \\ k+1 \end{pmatrix}$

$|\vec{AB}| = \sqrt{(k-2)^2 + (k+2)^2} = \sqrt{2k^2 + 8}$

$|\vec{AD}| = \sqrt{(k-3)^2 + (k+1)^2} = \sqrt{2k^2 - 4k + 10}$

Equating and squaring both sides: $2k^2 + 8 = 2k^2 - 4k + 10$

$4k = 2$

$k = \frac{1}{2}$

Video: [Vector geometry A](#)

Video: [Vector geometry B](#)

[Solutions to Starter and E.g.s](#)

Exercise

p242 12D Qu 2i, 3iabc, 4-7, 9-11

